

Multiple Sclerosis Enrollment Form

330	2 0.5. Highway 5. Howell. N.	.5. 077511 www. pai kw	PATIENT INFORMATION	III FITOTIE: 1-866-355-7797 Fax. 1-88	00-001-0200	
Name: Date of Birth:				O Male O Female	O Male O Female	
Address:			City: State:	7in·	Zip:	
			; ;			
Phone: Alt Phone: SS #: Primary Language:				Email:		
SS #:		Emergency Contact:	Emergency Contact:			
PRESCRIBER INFORMATION						
Prescribing Practiti Supervising Physici		NPI#:	NPI#: NPI#:			
Address: City: State: Zip:				Tax ID:	IaxiD:	
Phone: Fax:			Office Contact:	Office Contact:		
			MEDICAL INFORMATION			
	** PLEASE FAX COPY OF PRESC	RIPTION MEDICATION/N	MEDICAL CARD, FRONT AND BACK, AS WELL Is patient new to therapy: O Yes O		IERAPY **	
O G35 Multiple Scl				No		
Date of Diagnosis:		first demyelinating ever				
Type:	O Relapsing-Remitting	O Secondary progressive with relapses			O Primary Progressive	
O Secondary progressive withou		ut relapses O Clinically Isolated Syndrome (CIS)			O Progressive-relapsing	
Drug Allergies: O NKDA						
PREVIOUS THERAPIES:		Strength and Dose Date of Therapy		keason for discontinuii	Reason for discontinuing:	
0		0				
0		0				
0		0	PRESCRIPTION INFORMATION			
Needs by Date:		Ship to: O Patients h	ome O Prescriber 1st order only O Prescr	riber all orders O Other		
Drug	Dose	Direction		Quantity	Refills	
-	O 30 mcg Pre-filled Syringe			O 4 week supply - 1 kit	(
O Avonex	O 30 mcg Single Dose Vial	O Inject 30 mcg intramuscularly once a week		O 12 week supply - 3 kits		
	O 30 Avonex Pen (single Dose)					
O Betaseron	O 0.3 mg	O Inject 0.25 mg (1ml) s	ubcutaneously every other day	O 28 day supply		
		O Dose Titration:		1 kit of 14 vials		
		Weeks 1-2: 0.0625 mg	(0.25ml) subcutaneously every other day			
		Weeks 3-4: 0.125 mg (0.50 ml) subcutaneously every other day	O 84 day supply		
		Weeks 5-6: 0.1875 mg	(0.75ml) subcutaneously every other day	3 kits of 14 vials		
		Weeks 7+: 0.25 mg (1m	l) subcutaneously every other day			
		O Other:		O Other:		
	O BETAJECT Lite Autoinjector	O Use as directed		0		
O Copaxone	O 20 mg Pre-filled Syringe	O Inject 20 mg subcutaneously daily		O 30 day supply		
				O 90 day supply		
	O Autoject 2			O 1 Box - 28 capsules		
O Gilenya	O 0.5 mg	O Take one capsule by mouth once daily		O 1 Box - 28 capsules		
		O Other:				
O Rebif O Rebif Redidose	O 8.8 mcg (0.2ml)	1	utaneously three times weekly for weeks 1-2,		O 4 week supply	
	O 22 mcg (0.5ml)	1	ıbcutaneously three times weekly for weeks 3	O 4 week supply		
	O 44 mcg (0.5ml)		utaneously three times weekly			
		O 44 mcg (0.5ml) subcutaneously three times weekly O Inject 0.25mg (1ml) subcutaneously every other day				
O Extavia			ubcutaneously every other day	O 30 day supply - 1 kit		
		O Dose Titraion:		O 90 day supply - 3 kits		
		Weeks 1-2: 0.0625mg	(0.25ml) subcutaneously every other day			
	O 0.3 mg	Weeks 3-4: 0.125mg (0	.50ml) subcutaneously every other day			
		Weeks 5-6: 0.1875mg	(0.75ml) subcutaneously every other day			
		:				
		Weeks 7+: 0.25 mg (1ml) subcutaneously every other day				
	O FVTAVIA Auto Injector II	O Other:		 0		
O EXTAVIA Auto Injector II O Use as directed O O					<u>:</u>	
O Patient has received pen and injection training O Physician's office to provide injection training O Parkway Pharmacy to coordinate injection training						
		Р	RESCRIBING PRACTITIONER SIGNATURE			
			ou are also authorizing Parkway Pharmacy to s	serve as your prior authorization designate	ed agent in dealing with	
medical and prescr	ription insurance companies, and	co-pay assistance found	dations.			
Droccribina Dro	tionor:			Dato		
Prescribing Practi	tioner.		CONFIDENTIALITY NOTICE	Date		
IMPORTANT: This fax is	intended to be delivered only to the nar	med addressee. It contains ma	sterial that is confidential, proprietary or exempt from disc	dosure under applicable law. If you are not the name	ed addressee, you should not	
			ived this document in error and then destroy this docume			
Faxed Prescription	s will only be accepted from a pre	escribing practitioner.				